

RECURSION OPERATORS OF SOME EQUATIONS OF HYDRODYNAMIC TYPE

M. Gürses and K. Zheltukhin

Department of Mathematics, Faculty of Sciences

Bilkent University, 06533 Ankara-Turkey

email: gurses@fen.bilkent.edu.tr

February 8, 2008

Abstract

We give a general method for constructing recursion operators for some equations of hydrodynamic type, admitting a nonstandard Lax representation. We give several examples for $N = 2$ and $N = 3$ containing the equations of shallow water waves and its generalizations with their first two higher symmetries and their recursion operators. We also discuss a reduction of $N + 1$ systems to N systems of some new equations of hydrodynamic type.

I. Introduction

Most of the integrable nonlinear partial differential equations admit Lax representations

$$L_t = [A, L], \quad (1)$$

where L is a pseudo-differential operator of order m and A is a pseudo differential operator. Recently [1] we established a new method for such integrable equations to construct their recursion operators. This method uses the hierarchy of equations

$$L_{t_n} = [A_n, L] \quad (2)$$

and the Gel'fand-Dikii [2] construction of the A_n -operators. Defining an operator R_n in the form

$$A_n = LA_{n-m} + R_n, \quad (3)$$

one then obtains relations among the hierarchies

$$L_{t_n} = LL_{t_{n-m}} + [R_n; L] \quad (4)$$

This equation allows to find L_{t_n} in terms of $L_{t_{n-m}}$. It is important to note that one does not need to know the exact form of A_n . For further details of the method see [1].

Here we extend this method to equations of hydrodynamic type [3]. These equations and their Hamiltonian formulation (sometimes called the dispersion-less KdV system) were studied by Dubrovin and Novikov [4]. See [5] for more details on this subject. It is known that these equations admit a nonstandard Lax representation

$$\frac{\partial L}{\partial t} = \{A, L\}_k, \quad (5)$$

where A, L are differentiable functions of t, x, p on a Poisson manifold M with local coordinates (x, p) and $\{, \}_k$ is the Poisson bracket. On M we take this Poisson bracket $\{, \}_k = p^k \{, \}$, where $\{, \}$ is the canonical Poisson bracket and k is an integer. For more information on Poisson manifolds see [6],[7]. Equations of hydrodynamic type with the above Lax representations were

studied in [8]-[11]. Having such Lax representation, we can consider a whole hierarchy of equations

$$\frac{\partial L}{\partial t_n} = \{A_n, L\}_k. \quad (6)$$

We can also represent function A_{n+m} in the form given in (3) and apply our method [1] for constructing of a recursion operator for the equation (6). There are some other works [12]- [14] which give also recursion operators of some equations of hydrodynamic type. The form of these operators are different then the recursion operators presented in this work. Our method [1] produces recursion operators for hydrodynamic type of equations in the form $\mathcal{R} = A + B D^{-1}$ where A and B are functions of dynamical variables and their derivatives. All higher symmetries obtained by the repeated application of this recursion operator to translational symmetries belong also to the hydrodynamic type of equations. The recursion operators obtained in references [12]-[14] are of the form $\mathcal{R} = C D + A + B D^{-1} E$, where A, B, C , and E are functions of dynamical variables and their derivatives.

In the next section we discuss the Lax representation with Poisson brackets for polynomial Lax functions. In Sec.III we give the method of construction of the recursion operators following [1]. In Sec.IV we give several examples for $k = 0$ and $k = 1$. In Sec.V we consider the Poisson bracket for general k and let

$$L = p + S + Pp^{-1}, \quad (7)$$

and find the Lax equations and the corresponding recursion operator for $N = 2$. In Sec.VI we consider the Lax function

$$L = p^{\gamma-1} + u + \frac{v^{\gamma-1}}{(\gamma-1)^2} p^{-\gamma+1}, \quad (8)$$

and take $k = 0$. We obtain the equations corresponding to the polytropic gas dynamics and its recursion operators [6], [10]. It is interesting to note that the systems of equations and their recursion operators obtained in these sections V and VI are transformable into each other. In Sec.VII we give a method reduction from $N + 1$ system to an N system by letting one of the symmetrical variables (defined in the text) to zero. Reduced systems are shown to be also integrable, i.e., they admit recursion operators.

II. Lax formulation with Poisson bracket

We start with the definition of the standard Poisson bracket. Let $f(x, p)$ and $g(x, p)$ be differentiable functions of their arguments. Then the standard Poisson bracket is defined by (see [6] and [9] for more details)

$$\{f, g\} = \frac{\partial f}{\partial p} \frac{\partial g}{\partial x} - \frac{\partial f}{\partial x} \frac{\partial g}{\partial p}. \quad (9)$$

We give a slight modification of this bracket as [9]

$$\{f, h\}_k = p^k \{f, g\}, \quad (10)$$

where k is an integer. Then we have

Lemma 1. *For any $k \in \mathbb{Z}$ the bracket $\{f, g\}_k$ defines a Poisson brackets.*

Proof. We should check only the Jacobi identity. Other properties of Poisson brackets are evidently true. The standard bracket $\{f, g\}$ satisfies the Jacobi identity. For all k we have to show that

$$\{\{f, g\}_k, h\}_k + \{\{h, f\}_k, g\}_k + \{\{g, h\}_k, f\}_k = 0.$$

First, note that

$$\{\{f, g\}_k, h\}_k = p^k \{p^k \{f, g\}, h\} = p^{2k} \{\{f, g\}, h\} + kp^{k-1} \{f, g\} h_x.$$

Thus, we have

$$\begin{aligned} \{\{f, g\}_k, h\}_k + \{\{h, f\}_k, g\}_k + \{\{g, h\}_k, f\}_k &= p^{2k} (\{\{f, g\}, h\} + \{\{h, f\}, g\} + \\ &\quad \{\{g, h\}, f\}) + kp^{k-1} (\{f, g\} h_x + \{h, f\} g_x + \{g, h\} f_x). \end{aligned}$$

Equality

$$\{\{f, g\}, h\} + \{\{h, f\}, g\} + \{\{g, h\}, f\} = 0$$

holds and it is easy to check that

$$\{f, g\} h_x + \{h, f\} g_x + \{g, h\} f_x = 0.$$

Hence, $\{, \}_k$, for all k defines a Poisson bracket. $\square$

For each $k \in \mathbf{Z}$ we can consider hierarchies of equations of hydrodynamic type, defined in terms of the Lax function

$$L = p^{N-1} + \sum_{i=-1}^{N-2} p^i S_i(x, t) \quad (11)$$

by the Lax equation

$$\frac{\partial L}{\partial t_n} = \left\{ (L^{\frac{n}{N-1}})_{\geq -k+1}; L \right\}_k, \quad (12)$$

where $n = j + l(N-1)$ and $j = 1, 2, \dots, (N-1)$, $l \in \mathbf{N}$. So we have a hierarchy for each k and $j = 1, \dots, (N-1)$. Also, we require $n \geq -k+1$ to ensure that $(L^{\frac{n}{N-1}})_{\geq -k+1}$ is not zero. With the choice of Poisson brackets $\{, \}_k$, we must take a certain part of the series expansion of $L^{\frac{n}{N-1}}$ to get the consistent equation (12). This part is $(L^{\frac{n}{N-1}})_{\geq -k+1}$.

The Lax function (11) can also be written in terms of symmetric variables $u_1, \dots, u_N$

$$L = \frac{1}{p} \prod_{j=1}^N (p - u_j) \quad (13)$$

that is $u_1, \dots, u_N$ are roots of the polynomial

$$p^{N-1} + S_{N-2}p^{N-2} + \dots + S_{-1}.$$

In new variables the equation (12) is invariant under transposition of variables.

III. Recursion Operators

For each hierarchy of the equations (12), depending on the pair (N, k) , we can find a recursion operator.

Lemma 2. *For any n*

$$L_n = LL_{n-(N-1)} + \{R_n; L\}_k, \quad (14)$$

where function R_n has a form

$$R_n = \sum_{i=0}^{N-2} p^{i-k} A_i(S_{-1} \dots S_{N-2}, S_{-1,n-1} \dots S_{N-2,n-1}). \quad (15)$$

Proof.

$$(L^{\frac{n}{N-1}})_{\geq -k+1} = [L(L^{\frac{n}{N-1}-1})_{\geq -k+1} + L(L^{\frac{n}{N-1}-1})_{< -k+1}]_{\geq -k+1}$$

So,

$$(L^{\frac{n}{N-1}})_{\geq -k+1} = L(L^{\frac{n}{N-1}-1})_{\geq -k+1} + (L(L^{\frac{n}{N-1}-1})_{< -k+1})_{\geq -k+1} - (L(L^{\frac{n}{N-1}-1})_{\geq -k+1})_{< -k+1}. \quad (16)$$

If we put

$$R_n = (L(L^{\frac{n}{N-1}-1})_{< -k+1})_{\geq -k+1} - (L(L^{\frac{n}{N-1}-1})_{\geq -k+1})_{< -k+1} \quad ,$$

then

$$(L^{\frac{n}{N-1}})_{\geq -k+1} = L(L^{\frac{n}{N-1}-1})_{\geq -k+1} + R_n.$$

Hence,

$$\begin{aligned} L_n &= \left\{ (L^{\frac{n}{N-1}})_{\geq -k+1}; L \right\}_k = \left\{ L(L^{\frac{n}{N-1}-1})_{\geq -k+1} + R_n; L \right\}_k \\ &= LL_{n-(N-1)} + \{R_n; L\}_k, \end{aligned} \quad (17)$$

and (14) is satisfied. Evaluating powers of $(L(L^{\frac{n}{N-1}-1})_{< -k+1})_{\geq -k+1}$ and $-(L(L^{\frac{n}{N-1}-1})_{\geq -k+1})_{< -k+1}$ we get that R_n has form (15). $\square$

Lemma 3. *A recursion operator for the hierarchy (12) is given by equalities, for $m = N-2, N-3, \dots, -1$,*

$$\begin{aligned} S_{m,n+(N-1)} &= \sum_{j=-1}^{m+1} S_j S_{m-j,n} + \sum_{j=-1}^{m+1} (j+1-k) A_{j+1} S_{m-j,x} - \\ &\quad \sum_{j=-1}^{m+1} (m-j) A_{j+1,x} S_{m-j}, \end{aligned} \quad (18)$$

where to simplify the above formula we have defined that $S_{N-1} = 1$ and $S_{N-1,x} = 0$, $S_{N-1,n} = 0$. Coefficients $A_{N-2}, A_{N-3}, \dots, A_0$ can be found from the recursion relations, for $m = N-2, \dots, -1$

$$(N-1)A_{m,x} = \sum_{j=m}^{N-1} S_j S_{N-2+m-j} + \sum_{j=m}^{N-2} (j+1-k) A_{j+1} S_{N-2+m-j,x} - \sum_{j=m}^{N-2} (N-2+m-j) A_{j+1,x} S_{N-2+m-j} \quad (19)$$

Proof. Let us write the equality (14), using (15) for R_n

$$\begin{aligned} \sum_{i=-1}^{N-2} p^i S_{i,n+(N-1)} &= \left(p^{N-1} + \sum_{i=-1}^{N-2} p^i S_i \right) \left(\sum_{i=-1}^{N-2} p^i S_{i,n} \right) + \\ &\quad p^k \left(\sum_{j=0}^{N-1} (j-k) p^{j-k-1} A_j \right) \left(\sum_{j=-1}^{N-2} p^j S_{j,x} \right) \\ &\quad - p^k \left(\sum_{j=0}^{N-1} p^{j-k} A_{j,x} \right) \left((N-1) p^{N-2} + \sum_{j=-1}^{N-2} j p^{j-1} S_j \right) \end{aligned}$$

To have the equality, the coefficients of $p^{2N-3}, \dots, p^{N-1}$ and p^{-2} must be zero, it gives recursion relations to find $A_{N-2}, \dots, A_0$. The coefficients of $p^{N-2}, \dots, p^{-1}$ give the expressions for $S_{N-2,n+(N-1)}, \dots, S_{-1,n+(N-1)}$. $\square$

Although the recursion operator $\mathcal{R}$, given by (18), is a pseudo-differential operator, but it gives a hierarchy of local symmetries starting from the equation itself. Indeed, equalities (18), (19) give expressions $S_{N-2,n+(N-1)}, \dots, S_{-1,n+(N-1)}$ in terms of $S_{N-2}, \dots, S_{-1}$ and $S_{N-2,n}, \dots, S_{-1,n}$. Hence, the recursion operator $\mathcal{R}$ is constructed in such a way that

$$\left\{ (L^{\frac{n}{N-1}+1})_{\geq -k+1}; L \right\}_k = \mathcal{R} \left(\left\{ (L^{\frac{n}{N-1}})_{\geq -k+1}; L \right\}_k \right) \quad (20)$$

IV. Some Integrable Systems

We shall consider first some examples for $k = 0$, $k = 1$ and the general case in the next section.

A. Multicomponent hierarchy containing also the shallow water wave equations , $k = 0$

This hierarchy corresponds to the case $k = 0$. Let us give the first equation of hierarchy and a recursion operator for $N = 2, 3$.

Proposition 1. *In the case $N = 2$ one has the Lax function*

$$L = p + S + P p^{-1}$$

and the Lax equation for $n = 2$, given by (47), when $k = 0$

$$\begin{aligned} \frac{1}{2}S_t &= SS_x + P_x, \\ \frac{1}{2}P_t &= SP_x + PS_x. \end{aligned} \tag{21}$$

and the recursion operator, given by (48),

$$\mathcal{R} = \begin{pmatrix} S + S_x D_x^{-1} & 2 \\ 2P + P_x D_x^{-1} & S \end{pmatrix}. \tag{22}$$

These equations are known as the shallow water wave equations or as the equations of polytropic gas dynamics for $\gamma = 2$ (See Sec.VI).

The first two symmetries of the system (21) are given by

$$\begin{aligned} S_{t_1} &= (S^3 + 6SP)_x, \\ P_{t_1} &= (3S^2P + 3P^2)_x, \end{aligned} \tag{23}$$

$$\begin{aligned} S_{t_2} &= (S^4 + 12S^2P + 6P^2)_x, \\ P_{t_2} &= (4S^3P + 12SP^2)_x. \end{aligned} \tag{24}$$

These are all commuting symmetries.

Remark 1. In symmetric variables the system (21) is written as

$$\begin{aligned}\frac{1}{2}u_t &= (u+v)u_x + uv_x, \\ \frac{1}{2}v_t &= vu_x + (u+v)v_x,\end{aligned}\tag{25}$$

and the recursion operator (22) takes the form

$$\mathcal{R} = \begin{pmatrix} u+v+u_xD_x^{-1} & 2u+u_xD_x^{-1} \\ 2v+v_xD_x^{-1} & u+v+v_xD_x^{-1} \end{pmatrix}.\tag{26}$$

Proposition 2. In the case $N = 3$ one has the Lax function

$$L = p^2 + pS + P + p^{-1}Q$$

and the Lax equation with $n = 3$ is

$$\begin{aligned}\frac{1}{3}S_t &= (\frac{1}{2}P - \frac{1}{8}S^2)S_x + \frac{1}{2}SP_x + Q_x, \\ \frac{1}{3}P_t &= \frac{1}{2}QS_x + (\frac{1}{8}S^2 + \frac{1}{2}P)P_x + SQ_x, \\ \frac{1}{3}Q_t &= \frac{1}{4}SQS_x + \frac{1}{2}QP_x + (\frac{1}{8}S^2 + \frac{1}{2}P)Q_x.\end{aligned}\tag{27}$$

The recursion operator, corresponding to this equation, is

$$\mathcal{R} = \begin{pmatrix} -\frac{S^2}{4} + P + P_xD_x^{-1} - \frac{S_x}{4}D_x^{-1} \cdot S & \frac{S}{2} + \frac{S_x}{2}D_x^{-1} & 3 \\ \frac{3Q}{2} + (Q_x + \frac{P_xS}{2})D_x^{-1} - \frac{P_x}{4}D_x^{-1} \cdot S & P + \frac{P_x}{2}D_x^{-1} & 2S \\ \frac{SQ}{4} + (\frac{SQ_x}{2} + \frac{S_xQ}{2})D_x^{-1} - \frac{Q_x}{4}D_x^{-1} \cdot S & \frac{3Q}{2} + \frac{Q_x}{2}D_x^{-1} & P \end{pmatrix}.\tag{28}$$

Proof. Using (19) we find the function R_n and using (18) we find the recursion operator (28). $\square$

Remark 2. In symmetric variables the equation (27) is written as

$$\begin{aligned}\frac{1}{3}u_t &= (-\frac{1}{8}u^2 + \frac{1}{2}(uv + uw + vw) + \frac{1}{8}(v+w)^2)u_x \\ &\quad + (\frac{1}{4}u^2 + \frac{1}{4}uv + \frac{3}{4}uw)v_x + (\frac{1}{4}u^2 + \frac{1}{4}uw + \frac{3}{4}uv)w_x, \\ \frac{1}{3}v_t &= (\frac{1}{4}v^2 + \frac{1}{4}uv + \frac{3}{4}vw)u_x + (\frac{1}{4}v^2 + \frac{1}{4}vw + \frac{3}{4}uv)w_x \\ &\quad + (-\frac{1}{8}v^2 + \frac{1}{2}(uv + uw + vw) + \frac{1}{8}(u+w)^2)v_x, \\ \frac{1}{3}w_t &= (\frac{1}{4}w^2 + \frac{1}{4}uw + \frac{3}{4}wv)u_x + (\frac{1}{4}w^2 + \frac{1}{4}wv + \frac{3}{4}uw)v_x \\ &\quad + (-\frac{1}{8}w^2 + \frac{1}{2}(uv + uw + vw) + \frac{1}{8}(v+u)^2)w_x,\end{aligned}\tag{29}$$

and the recursion operator takes the form (67) given in the Appendix.

B. Toda hierarchy ($k = 1$)

Toda hierarchy corresponds to the case $k = 1$ [9]. Let us give the first equation of hierarchy and a recursion operator for $N = 2$ and $N = 3$.

Proposition 3. *In the case $N = 2$ and $n = 1$ one has the Lax function*

$$L = p + S + P p^{-1}$$

and the Lax equation for $n = 1$, given by (41),

$$\begin{aligned} S_t &= P_x, \\ P_t &= P S_x, \end{aligned} \tag{30}$$

and the recursion operator, given by (42),

$$\mathcal{R} = \begin{pmatrix} S & 2 + P_x D_x^{-1} \cdot P^{-1} \\ 2P & S + S_x P D_x^{-1} \cdot P^{-1} \end{pmatrix}. \tag{31}$$

The first two symmetries of the equation (30) are given by

$$\begin{aligned} S_{t_1} &= (2SP)_x, \\ P_{t_1} &= P(2P + S^2)_x, \end{aligned} \tag{32}$$

$$\begin{aligned} S_{t_2} &= (3S^2P + 3P^2)_x, \\ P_{t_2} &= P(6PS + S^3)_x. \end{aligned} \tag{33}$$

Remark 3. *In symmetric variables the equation (30) is written as*

$$\begin{aligned} u_t &= uv_x, \\ v_t &= vu_x, \end{aligned} \tag{34}$$

and the recursion operator (31) takes the form

$$\mathcal{R} = \begin{pmatrix} u + v + uv_x D_x^{-1} \cdot u^{-1} & 2u + uv_x D_x^{-1} \cdot v^{-1} \\ 2v + vu_x D_x^{-1} \cdot u^{-1} & u + v + vu_x D_x^{-1} \cdot v^{-1} \end{pmatrix}. \tag{35}$$

Proposition 4. *In the case $N = 3$ and $n = 1$ one has the Lax function*

$$L = p^2 + pS_1 + P + p^{-1}Q$$

and the Lax equation with $n = 1$ is

$$\begin{aligned} S_t &= P_x - \frac{1}{2}SS_x, \\ P_t &= Q_x, \\ Q_t &= \frac{1}{2}QS_x. \end{aligned} \tag{36}$$

The recursion operator, corresponding to this equation, is

$$\mathcal{R} = \begin{pmatrix} P - \frac{1}{4}S^2 + (\frac{1}{2}P_x - \frac{1}{4}SS_x)D_x^{-1} & \frac{1}{2}S & 3 + 2Q_xD_x^{-1} \cdot Q^{-1} \\ \frac{3}{2}Q + \frac{1}{2}Q_xD_x^{-1} & P & 2S + (SQ)_xD_x^{-1} \cdot Q^{-1} \\ \frac{1}{4}SQ + \frac{1}{4}S_xQD_x^{-1} & \frac{3}{2}Q & P + P_xQD_x^{-1} \cdot Q^{-1} \end{pmatrix}. \tag{37}$$

Proof. Using equalities (19) we find the function R_n and using (18) we find the recursion operator (37). $\square$

Remark 4. *In symmetric variables the equation (36) is written as*

$$\begin{aligned} u_t &= \frac{1}{2}u(-u_x + v_x + w_x), \\ v_t &= \frac{1}{2}v(+u_x - v_x + w_x), \\ w_t &= \frac{1}{2}w(+u_x + v_x - w_x), \end{aligned} \tag{38}$$

and the recursion operator takes the form (68) given in the Appendix.

V. Lax equation for general k

We shall only consider the case where $N = 2$. We have the Lax function

$$L = p + S + Pp^{-1} \tag{39}$$

and the Lax equation

$$\frac{\partial L}{\partial t_n} = \{(L^n)_{\geq -k+1}; L\}_k. \tag{40}$$

We consider two cases $k \geq 1$ and $k \leq 0$.

A. The first case $k \geq 1$

Proposition 5. *In the case $N = 2$ and $k \geq 1$ one has the Lax equation*

$$\begin{aligned} S_t &= kP^{k-1}P_x, \\ P_t &= kP^kS_x. \end{aligned} \quad (41)$$

and the recursion operator for this equation is

$$\mathcal{R} = \begin{pmatrix} S + (1-k)S_xD_x^{-1} & 2 + kP^{k-1}P_xD_x^{-1} \cdot P^{-k} \\ 2P + (1-k)P_xD_x^{-1} & S + kS_xP^kD_x^{-1} \cdot P^{-k} \end{pmatrix}. \quad (42)$$

Proof. The smallest power of p in L^n is $-n$. To have powers less than $-k+1$ we must put $n = k$. If there are no such powers then Poisson brackets are $\{(L^n); L\}_k = 0$.

Let us calculate the Lax equation

$$L_t = \{(L^k)_{\geq -k+1}; L\}_k = -\{(L^k)_{\leq -k}; L\}_k$$

We have $(L^k)_{\leq -k} = [(p + S + Pp^{-1})^k]_{\leq -k} = P^k p^{-k}$, thus

$$L_t = -\{P^k p^{-k}; p + S + Pp^{-1}\}_k.$$

And we get the equation (41). Using (18), (19) we find the recursion operator (42). $\square$

First two symmetries are given as follows

$$\begin{aligned} S_{t_1} &= (P^k S)_x, \\ P_{t_1} &= P^k \left(P + \frac{k}{2} S^2 \right)_x, \end{aligned} \quad (43)$$

$$\begin{aligned} S_{t_2} &= (k+1)(k+2) \left(\frac{1}{2} P^k S^2 + \frac{1}{k+1} P^{k+1} \right)_x, \\ P_{t_2} &= (k+1)(k+2) P^k \left(PS + \frac{k}{6} S^3 \right)_x. \end{aligned} \quad (44)$$

Remark 5. *In symmetric variables the equation (41) is written as*

$$\begin{aligned} u_t &= ku^k v^{k-1} v_x, \\ v_t &= ku^{k-1} v^k u_x, \end{aligned} \quad (45)$$

and the recursion operator (42) takes the form

$$\mathcal{R} = \begin{pmatrix} u + v + (1-k)u_x D_x^{-1} + & 2u + (1-k)u_x D_x^{-1} + \\ ku^k v^{k-1} v_x D_x^{-1} \cdot u^{-k} v^{-k+1} & ku^k v^{k-1} v_x D_x^{-1} \cdot u^{-k+1} v^{-k} \\ \\ 2v + (1-k)v_x D_x^{-1} + & u + v + (1-k)v_x D_x^{-1} + \\ ku^{k-1} v^k u_x D_x^{-1} \cdot u^{-k} v^{-k+1} & ku^{k-1} v^k u_x D_x^{-1} \cdot u^{-k+1} v^{-k} \end{pmatrix}. \quad (46)$$

B. The second case $k \leq 0$

Proposition 6. *In the case $N = 2$ and $k \leq 0$ one has the Lax equation*

$$\begin{aligned} S_t &= (-k+2)(-k+1)SS_x + (-k+2)P_x, \\ P_t &= (-k+2)(-k+1)SP_x + (-k+2)S_x P. \end{aligned} \quad (47)$$

and the recursion operator for this equation is

$$\mathcal{R} = \begin{pmatrix} S + (1-k)S_x D_x^{-1} & 2 + kP^{k-1}P_x D_x^{-1} \cdot P^{-k} \\ 2P + (1-k)P_x D_x^{-1} & S + kS_x P^k D_x^{-1} \cdot P^{-k} \end{pmatrix}. \quad (48)$$

Proof. The largest power of p in L^n is p^n . To have powers larger than $-k+1$ we must put $n = -k+1$. Then we have

$$(L^{-k+1})_{\geq -k+1} = [(p + S + Pp^{-1})^{-k+1}]_{\geq -k+1} = p^{-k+1},$$

thus

$$L_t = \{p^{-k+1}; p + S + Pp^{-1}\}_k.$$

Then the Lax equation becomes

$$\begin{aligned} S_t &= S_x, \\ P_t &= P_x. \end{aligned}$$

This is a trivial equation, let us calculate the second symmetry. We have $(L^{-k+2})_{\geq -k+1} = [(p + S + Pp^{-1})^{-k+1}]_{\geq -k+1} = p^{-k+2} + (-k+2)Sp^{-k+1}$, thus

$$L_t = \{p^{-k+2} + (-k+2)Sp^{-k+1}; p + S + Pp^{-1}\}_k.$$

we get the equation (47). Using (18), (19) we find the recursion operator (48). $\square$

First two symmetries are given as follows

$$\begin{aligned} S_{t_1} &= (k-2)(k-3) \left(P S + \frac{1}{6}(1-k)S^3 \right)_x, \\ P_{t_1} &= (k-2)(k-3) \left(S S_x P + \frac{1}{2}(1-k)S^2 P_x + P P_x \right), \end{aligned} \quad (49)$$

$$\begin{aligned} S_{t_2} &= (2-k)(3-k)(4-k) \left(\frac{1}{2}S^2 P + \frac{1}{6}S^4 + \frac{1}{2(2-k)}P^2 \right)_x, \\ P_{t_2} &= (2-k)(3-k)(4-k) \left(\frac{1}{2}S^2 S_x P + \frac{1}{6}(1-k)S^3 P_x \right. \\ &\quad \left. + S P P_x + \frac{1}{(2-k)}P^2 S_x \right). \end{aligned} \quad (50)$$

Remark 6. In symmetric variables the equation (47) is written as

$$\begin{aligned} u_t &= (-k+2)(1-k)(u+v)u_x + (-k+2)uv_x, \\ v_t &= (-k+2)vu_x + (-k+2)(1-k)(u+v)v_x, \end{aligned} \quad (51)$$

and the recursion operator (48) takes the form

$$\mathcal{R} = \begin{pmatrix} u + v + (1-k)u_x D_x^{-1} + & 2u + (1-k)u_x D_x^{-1} + \\ k u^k v^{k-1} v_x D_x^{-1} \cdot u^{-k} v^{-k+1} & k u^k v^{k-1} v_x D_x^{-1} \cdot u^{-k+1} v^{-k} \\ \\ 2v + (1-k)v_x D_x^{-1} + & u + v + (1-k)v_x D_x^{-1} + \\ k u^{k-1} v^k u_x D_x^{-1} \cdot u^{-k} v^{-k+1} & k u^{k-1} v^k u_x D_x^{-1} \cdot u^{-k+1} v^{-k} \end{pmatrix}. \quad (52)$$

In this Section, to obtain the recursion operators we have considered two different cases $k \leq 0$ and $k \geq 1$ to simplify some technical problems in the method. At the end we obtained recursion operators having the same forms (42) and (48). Hence any one of these represent the recursion operator for $k \in \mathbb{Z}$. It seems, comparing the results, that the systems of equations in one case are symmetries of the other case. For instance, the system (47) is a symmetry of system (41). Hence we may consider only one case with recursion operator (42) for all integer values of k .

VI. Lax function for polytropic gas dynamics

In this section we consider another Lax function, introduced in [10],

$$L = p^{\gamma-1} + u + \frac{v^{\gamma-1}}{(\gamma-1)^2} p^{-\gamma+1} \quad (53)$$

and the Lax equation

$$\frac{\partial L}{\partial t} = \frac{\gamma-1}{\gamma} \left\{ (L^{\frac{\gamma}{\gamma-1}})_{\geq 1}, L \right\}_0, \quad (54)$$

gives the equations of the polytropic gas dynamics.

Proposition 7. *The Lax equation corresponding to (54) is*

$$\begin{aligned} u_t + uu_x + v^{\gamma-2}v_x &= 0, \\ v_t + (uv)_x &= 0. \end{aligned} \quad (55)$$

Proof. Expanding the function (53) around the point $p = \infty$, we have

$$\left(p^{\gamma-1} + u + \frac{v^{\gamma-1}}{(\gamma-1)^2} p^{-\gamma+1} \right)^{\frac{\gamma}{\gamma-1}} = p^\gamma + \frac{\gamma}{\gamma-1} pu + \dots$$

all other terms have negative powers of p . Therefore

$$\left(L^{\frac{\gamma}{\gamma-1}} \right)_{\geq 1} = p^\gamma + \frac{\gamma}{\gamma-1} pu$$

and the Lax equation (54) corresponds to (55). $\square$

Proposition 8. *The recursion operator for the equation (55) is*

$$\mathcal{R} = \begin{pmatrix} u + \frac{u_x}{\gamma-1} D_x^{-1} & \frac{2v^{\gamma-2}}{\gamma-1} + \frac{(v^{\gamma-2})_x}{\gamma-1} D_x^{-1} \\ \frac{2v}{\gamma-1} + \frac{v_x}{\gamma-1} D_x^{-1} & u + \frac{\gamma-2}{\gamma-1} u_x D_x^{-1} \end{pmatrix}. \quad (56)$$

Proof. Using the equation

$$\frac{\partial L}{\partial t_{n+1}} = L \frac{\partial L}{\partial t_n} + \{R_n, L\}.$$

in the same way as for polynomial Lax function one can find the recursion operator (56). $\square$

It is interesting to note that the equation (47) and equations of polytropic gas dynamics (55) are related by the following change of variables

$$\begin{aligned} S &= \frac{u}{(-k+2)(-k+1)}, \\ P &= \frac{v^{\frac{1}{-k+1}}}{(-k+2)^2}, \end{aligned} \quad (57)$$

where $\gamma = \frac{-k+2}{-k+1}$. We note that under this change of variables recursion operator (48) is mapped to the recursion operator (56).

VII. Reduction

We consider reduction of the equation (12), written in symmetric variables, by setting $u_1 = 0$. Let us write the equation (12) as

$$\Delta(u_N, \dots, u_1) = 0, \quad (58)$$

where Δ is a differential operator. Then

$$\Delta(u_N, \dots, u_1)|_{u_1=0} = \left(\frac{\tilde{\Delta}(u_N, \dots, u_2)}{0} \right) \quad (59)$$

where $\tilde{\Delta}$ is another differential operator. Indeed, following [8] for the Lax function $L = \frac{1}{p} \prod_{j=1}^N (p - u_j)$ we have

$$\frac{\partial L}{\partial t} = L \sum_{j=1}^N \frac{u_{j,t}}{p + u_j},$$

$$\frac{\partial L}{\partial x} = L \sum_{j=1}^N \frac{u_{j,x}}{p + u_j}$$

and

$$\frac{\partial L}{\partial p} = L \left(-\frac{1}{p} + \sum_{j=1}^N \frac{1}{p + u_j} \right).$$

Thus $u_{j,t} = \text{Res}_{p=-u_j} \{M, L\}_k$, where $M = (L^{\frac{n}{N-1}})_{\geq -k+1}$. The Lax equation (12) can be written as

$$\sum_{j=1}^N \frac{u_{j,t}}{p + u_j} = p^k M_p \sum_{j=1}^N \frac{u_{j,x}}{p + u_j} - p^k M_x \left(-\frac{1}{p} + \sum_{j=1}^N \frac{1}{p + u_j} \right).$$

Note, that $p^k M_x$ and $p^k M_p$ are polynomials. So, if we put $u_1 = 0$ and calculate residue of right hand side at $p = 0$ we get (59). A new equation

$$\tilde{\Delta}(u_N, \dots, u_2) = 0 \quad (60)$$

is also integrable and a recursion operator of this equation can be obtained as reduction of the recursion operator of the equation (58). Let $\mathcal{R}$ be the recursion operator of (58) given by lemma 3, then

$$\mathcal{R}|_{u_1=0} = \left(\frac{\tilde{R}}{0 \dots 0} \middle| \frac{*}{0} \right). \quad (61)$$

Indeed, we found the recursion operator using formula (14). This formula can be written as

$$\sum_{j=1}^N \frac{u_{j,t_n}}{p + u_j} = L L_{n-(N-1)} + p^k R_{n,p} \sum_{j=1}^N \frac{u_{j,x}}{p + u_j} - p^k R_{n,x} \left(-\frac{1}{p} + \sum_{j=1}^N \frac{1}{p + u_j} \right)$$

and in the same way as for reduction of (58) we have (61), note, that $p^k R_{n,x}$ and $p^k R_{n,p}$ are also polynomials.

Lemma 4. *The operator $\tilde{R}$ is a recursion operator of the equation (60).*

Proof. Equation (60) is an evolution equation, so, to prove that $\tilde{R}$ is a recursion operator we must prove that for any solution $(u_N, \dots, u_2)$ of (60) the following equality holds (see [6])

$$D_{\tilde{\Delta}} \tilde{R} = \tilde{R} D_{\tilde{\Delta}},$$

where $D_{\tilde{\Delta}}$ is a Frechet derivative of $\tilde{\Delta}$.

If $(u_N, \dots, u_2)$ is a solution of (60) then $(u_N, \dots, u_2, u_1 = 0)$ is a solution of (58) and for the solution $(u_N, \dots, u_2, u_1 = 0)$ we have

$$D_{\Delta} \mathcal{R} = \mathcal{R} D_{\Delta}. \quad (62)$$

Next

$$D_{\Delta}|_{u_1=0} = \left(\begin{array}{c|c} \tilde{D} & * \\ \hline 0 \dots 0 & * \end{array} \right)$$

and

$$\mathcal{R}|_{u_1=0} = \left(\begin{array}{c|c} \tilde{R} & * \\ \hline 0 \dots 0 & 0 \end{array} \right).$$

Hence by (62) we have $\tilde{D}\tilde{R} = \tilde{R}\tilde{D}$. Calculating Frechet derivative we take derivatives with respect to one variable, considering other variables as constants. Thus, to calculate $\tilde{D}$ we can put $u_1 = 0$ and differentiate with respect to other variables or we can first differentiate and then put $u_1 = 0$. It means that $\tilde{D} = D_{\tilde{\Delta}}$ and

$$D_{\tilde{\Delta}} \tilde{R} = \tilde{R} D_{\tilde{\Delta}}.$$

□

Let us consider reduction of systems, given by remark 2 and remark 4 and their recursion operators.

Proposition 9. *Putting $w = 0$ in (38) and (68) we obtain a new system*

$$\begin{aligned} u_t &= \frac{1}{2}u(-u_x + v_x), \\ v_t &= \frac{1}{2}v(+u_x - v_x), \end{aligned} \quad (63)$$

and its recursion operator

$$\mathcal{R} = \begin{pmatrix} -uv + \frac{u}{4}(u+v) & -\frac{u}{4}(u+v) \\ +\frac{u}{4}(u_x - v_x)D_x^{-1} & +\frac{u}{4}(u_x - v_x)D_x^{-1} \\ -\frac{v}{4}(u+v) & -uv + \frac{v}{4}(u+v) \\ +\frac{v}{4}(-u_x + v_x)D_x^{-1} & +\frac{v}{4}(-u_x + v_x)D_x^{-1} \end{pmatrix} \quad (64)$$

respectively. $\square$

Proposition 10. Putting $w = 0$ in (29) and (67) we obtain a new system

$$\begin{aligned} \frac{1}{3}u_t &= (-\frac{1}{8}u^2 + \frac{1}{2}uv + \frac{1}{8}v^2)u_x + (\frac{1}{4}u^2 + \frac{1}{4}uv)v_x, \\ \frac{1}{3}v_t &= (\frac{1}{4}v^2 + \frac{1}{4}uv)u_x + (-\frac{1}{8}v^2 + \frac{1}{2}uv + \frac{1}{8}u^2)v_x, \end{aligned} \quad (65)$$

and its recursion operator

$$\mathcal{R} = \begin{pmatrix} -\frac{u^2}{4} + \frac{3uv}{4} + (\frac{u_x v}{2} + \frac{u v_x}{2})D_x^{-1} & \frac{u}{4}(u+v) + (\frac{u_x v}{2} + \frac{u v_x}{2})D_x^{-1} \\ -\frac{u_x}{4}D_x^{-1} \cdot u + \frac{u_x}{4}D_x^{-1} \cdot v & +\frac{u_x}{4}D_x^{-1} \cdot u - \frac{u_x}{4}D_x^{-1} \cdot v \\ \frac{v}{4}(u+v) + (\frac{u v_x}{2} + \frac{u_x v}{2})D_x^{-1} & -\frac{v^2}{4} + \frac{3uv}{4} + (\frac{u v_x}{2} + \frac{u_x v}{2})D_x^{-1} \\ -\frac{v_x}{4}D_x^{-1} \cdot u + \frac{v_x}{4}D_x^{-1} \cdot v & +\frac{v_x}{4}D_x^{-1} \cdot u - \frac{v_x}{4}D_x^{-1} \cdot v \end{pmatrix}, \quad (66)$$

respectively. $\square$

It is worth to mention that by reduction we obtain a new equation. For example, consider the case $k = 0$. The equation (25, corresponding to $N = 2$, and reduction of the equation (29), corresponding to $N = 3$, are not related by a linear transformation of variables. Indeed, in the equation (25)) coefficients of u_x, v_x are linear in u, v but in the equation (65) coefficients of u_x, v_x contain quadratic terms. Hence they can not be related by a linear transformation.

VII. Conclusion

We have constructed the recursion operators of some equations of hydrodynamic type. The form of these operators fall into the class of pseudo differential operators $A + B D^{-1}$ where A and B are functions of dynamical variables and their derivatives. The generalized symmetries of these equations are local and all belong to the same class (i.e., they are also equations

of hydrodynamic type). We have introduced a method of reduction which leads also to integrable class. These properties, bi Hamiltonian structure of the equations we obtained and equations with rational Lax functions will be communicated elsewhere.

Acknowledgments

We thank Burak Gürel and Atalay Karasu for several discussions. This work is partially supported by the Scientific and Technical Research Council of Turkey and by Turkish Academy of Sciences.

Appendix. Recursion operators of the systems (29) and (38) are respectively given by

$$\mathcal{R} = \begin{pmatrix} -\frac{u^2}{4} + \frac{3}{4}(uv + uw) + uv & \frac{u}{4}(u + v + w) + \frac{3uw}{2} & \frac{u}{4}(u + v + w) + \frac{3uv}{2} \\ +\frac{u_x}{2}(v + w)D_x^{-1} & +\frac{u_x}{2}(v + w)D_x^{-1} & +\frac{u_x}{2}(v + w)D_x^{-1} \\ +\frac{u}{2}(v_x + w_x)D_x^{-1} & +\frac{u}{2}(v_x + w_x)D_x^{-1} & +\frac{u}{2}(v_x + w_x)D_x^{-1} \\ -\frac{u_x}{4}D_x^{-1} \cdot u + \frac{u_x}{4}D_x^{-1} \cdot v & +\frac{u_x}{4}D_x^{-1} \cdot u - \frac{u_x}{4}D_x^{-1} \cdot v & +\frac{u_x}{4}D_x^{-1} \cdot u + \frac{u_x}{4}D_x^{-1} \cdot v \\ +\frac{u_x}{4}D_x^{-1} \cdot w & +\frac{u_x}{4}D_x^{-1} \cdot w & -\frac{u_x}{4}D_x^{-1} \cdot w \\ \\ \frac{v}{4}(u + v + w) + \frac{3vw}{2} & -\frac{v^2}{4} + \frac{3}{4}(uv + vw) + uv & \frac{v}{4}(u + v + w) + \frac{3uv}{2} \\ +\frac{v_x}{2}(u + w)D_x^{-1} & +\frac{v_x}{2}(u + w)D_x^{-1} & +\frac{v_x}{2}(u + w)D_x^{-1} \\ +\frac{v}{2}(u_x + w_x)D_x^{-1} & +\frac{v}{2}(u_x + w_x)D_x^{-1} & +\frac{v}{2}(u_x + w_x)D_x^{-1} \\ -\frac{v_x}{4}D_x^{-1} \cdot u + \frac{v_x}{4}D_x^{-1} \cdot v & +\frac{v_x}{4}D_x^{-1} \cdot u - \frac{v_x}{4}D_x^{-1} \cdot v & +\frac{v_x}{4}D_x^{-1} \cdot u + \frac{v_x}{4}D_x^{-1} \cdot v \\ +\frac{v_x}{4}D_x^{-1} \cdot w & +\frac{v_x}{4}D_x^{-1} \cdot w & -\frac{v_x}{4}D_x^{-1} \cdot w \\ \\ \frac{w}{4}(u + v + w) + \frac{3uw}{2} & \frac{w}{4}(u + v + w) + \frac{3uv}{2} & -\frac{w^2}{4} + \frac{3}{4}(uw + vw) + uv \\ +\frac{w_x}{2}(u + v)D_x^{-1} & +\frac{w_x}{2}(u + v)D_x^{-1} & +\frac{w_x}{2}(u + v)D_x^{-1} \\ +\frac{w}{2}(u_x + v_x)D_x^{-1} & +\frac{w}{2}(u_x + v_x)D_x^{-1} & +\frac{w}{2}(u_x + v_x)D_x^{-1} \\ -\frac{w_x}{4}D_x^{-1} \cdot u + \frac{w_x}{4}D_x^{-1} \cdot v & +\frac{w_x}{4}D_x^{-1} \cdot u - \frac{w_x}{4}D_x^{-1} \cdot v & +\frac{w_x}{4}D_x^{-1} \cdot u + \frac{w_x}{4}D_x^{-1} \cdot v \\ +\frac{w_x}{4}D_x^{-1} \cdot w & +\frac{w_x}{4}D_x^{-1} \cdot w & -\frac{w_x}{4}D_x^{-1} \cdot w \end{pmatrix}. \quad (67)$$

$$\mathcal{R} = \begin{pmatrix} \begin{array}{ccc} -\frac{u}{4}(uv + uw + vw) & -\frac{u}{4}(u + v + w) & -\frac{u}{4}(u + v + w) \\ +\frac{u}{4}(u + v + w) & -\frac{3uw}{2} & -\frac{3uv}{2} \\ +\frac{u}{4}(u_x - v_x - w_x)D_x^{-1} & +\frac{u}{4}(u_x - v_x - w_x)D_x^{-1} & +\frac{u}{4}(u_x - v_x - w_x)D_x^{-1} \\ -u(wv_x + vw_x)D_x^{-1} \cdot u^{-1} & -u(wv_x + vw_x)D_x^{-1} \cdot v^{-1} & -u(wv_x + vw_x)D_x^{-1} \cdot w^{-1} \end{array} \\ \begin{array}{ccc} -\frac{v}{4}(u + v + w) & -(uv + uw + vw) & -\frac{v}{4}(u + v + w) \\ -\frac{3vw}{2} & +\frac{v}{4}(u + v + w) & -\frac{3uv}{2} \\ +\frac{v}{4}(-u_x + v_x - w_x)D_x^{-1} & +\frac{v}{4}(-u_x + v_x - w_x)D_x^{-1} & +\frac{v}{4}(-u_x + v_x - w_x)D_x^{-1} \\ -v(wu_x + uw_x)D_x^{-1} \cdot u^{-1} & -v(wu_x + uw_x)D_x^{-1} \cdot v^{-1} & -v(wu_x + uw_x)D_x^{-1} \cdot w^{-1} \end{array} \\ \begin{array}{ccc} -\frac{w}{4}(u + v + w) & -\frac{w}{4}(u + v + w) & -(uv + uw + vw) \\ -\frac{3uw}{2} & -\frac{3vw}{2} & +\frac{w}{4}(u + v + w) \\ +\frac{w}{4}(-u_x - v_x + w_x)D_x^{-1} & +\frac{w}{4}(-u_x - v_x + w_x)D_x^{-1} & +\frac{w}{4}(-u_x - v_x + w_x)D_x^{-1} \\ -w(uv_x + vu_x)D_x^{-1} \cdot u^{-1} & -w(uv_x + vu_x)D_x^{-1} \cdot v^{-1} & -w(uv_x + vu_x)D_x^{-1} \cdot w^{-1} \end{array} \end{pmatrix}. \quad (68)$$

References

- [1] M. Gürses, A. Karasu, V.V. Sokolov, " On construction of recursion operator from Lax representation", *J. Math. Phys.*, **40**, 6473-6490 (1999).
- [2] I.M. Gel'fand, L.A. Dikii, Asymptotic Behavior of the Re-solvent of Sturm-Liouville equations and the Algebra of the Korteweg-de Vrise equations, *Funk. Anal. Appl.*, **10**, 13 (1976).
- [3] Preliminary version of this work was first reported in M. Gürses and K. Zhelthukin, "On Construction of Recursion Operators for Some Equations of Hydrodynamic Type", in *International Conference on Complex Analysis , Differential Equations and Related Topics*, May 29-June 3, 2000, Ufa, Russia.
- [4] B.A. Dubrovin and S.P. Novikov, "Hamiltonian formalism of one-dimensional systems of hydrodynamic type", *Soviet Math. Dokl.* , **27**, 665 (1983).

- [5] E.V. Ferepantov, "Hydrodynamic-type systems", in *CRC Handbook of Lie Group Analysis of Differential Equations*, Vol. 1. Ed. N.H. Ibragimov, p. 303-31. CRC Press, New York, 1994
- [6] P.J. Olver, *Applications of Lie Groups to Differential Equations*, Second Edition, Graduate Text in Mathematics, Vol. 107. Springer-Verlag, New-York (1993).
- [7] A. Weinstein, "Poisson Geometry", *Differential Geometry and its applications*, **9**, 213-238 (1998).
- [8] D.B. Fairlie and I.A.B. Strachan, "The algebraic and Hamiltonian structure of the dispersion-less Benney and Toda hierarchies", *Inverse Problems*, **12**, 885-908 (1998).
- [9] I.A.B. Strachan, "Degenerate Frobenius manifolds and the bi-Hamiltonian structure of rational Lax equations", *J. Math. Phys.*, **40**, 5058-5079 (1999).
- [10] J.C. Brunelli, A. Das, "A Lax description for polytropic gas dynamics", *Phys. Lett. A*, **235**, 597-602 (1997).
- [11] J.C. Brunelli, M. Gürses, and K. Zhelthukin, "On the integrability of some Monge-Ampere' equations", *Reviews in Mathematical Physics*, 2000.
- [12] M.B. Sheftel, "Generalized Hydrodynamic-type systems" in *CRC Handbook of Lie Group Analysis of Differential Equations*, Vol. 3. Edited by N.H. Ibragimov, p.169-189. CRC Press, New York, 1996.
- [13] V.M. Teshukov, "Hyperbolic systems admitting a nontrivial Lie-Backlund group", *LIIAN*, **106**, 25-30, 1989.
- [14] A.P. Fordy and B. Gürel, "A new construction of recursion operators for systems of hydrodynamic type", *Theoret. Math. Phys.*, 1999.